Pre-equilibrium Effects



- Issues
 - Surrogate technique relies on "same" compound nucleus
 - Parent reaction and "surrogate" reaction will have different PE
 - » Neutron reactions have 1p, 2p-1h, 3p-2h,....
 - » Surrogates more complex-e.g 0h, 1p-2h,2p-3h,...
 - Amount of PE emission will be reaction dependent
 - » HOW WELL CAN WE ESTIMATE PE EMISSION?
 - Phenomenological exiton models do pretty well
 - Residual Ang.Mom. will be reaction dependent?
 - » HOW WELL CAN WE ESTIMATE ANG. MOM.?
 - Much higher uncertainty
 - Quantum approaches may be more useful than semiclassical
 - For (n,γ) PE not important, BUT for surrogate PE could be large



Pre-equilibrium Effects




Discussion


- Lively several active participants
- Use extensive p,p' data inelastic and ang. Distributions
- Direct reaction data useful get higher accuracy with coupled channel analysis
- Reaction at 14 MeV different than 50 MeV look at knock out (quasi-elastic) reactions
- Surrogate studied nuclei may not have well known levels
- Experiments at GANIL on 90Zr good optical model fits
- Presentation on Kerman-McVoy approach for direct reactions formal mechanism for initial reaction stage -relevent to Younes/Britt - used with p-γ by Dietrich
- PE ang mom many approaches: FKK, NWY, TUL, and semiclassical models such as exciton and HMS.
- Exp GEANIE data 90 Zr



⁹⁰Zr(n, n' γ) / ⁹⁰Zr(n, n')







Shape (E_n > 14 meV) may reflect spin distribution in residual nucleus
GEANIE/WNR can resolve longstanding issues of preequilibrium



Pre-equilibrium Effects



Action Items

- Concentrate on 90Zr+n
 - » Calculate PE ang mom distribution vs neutron energy
 - » Finish analysis of GEANIE data for (n,n') gamma feeding
 - » GNASH/STAPRE/EMPIRE gamma emission from calculated PE ang mom distribution
 - » Compare against (n,xn)neutron emission spectra at 14, 18 MeV
 - » Compare Exp/Model to determine PE ang mom understanding
 - » Do similar comparisons with other data sets.

